

## PCT COOPERATION TREATY

## PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner  
 US Department of Commerce  
 United States Patent and Trademark  
 Office, PCT  
 2011 South Clark Place Room  
 CP2/5C24  
 Arlington, VA 22202  
 ETATS-UNIS D'AMERIQUE  
 in its capacity as elected Office

<b>Date of mailing (day/month/year)</b> 09 February 2001 (09.02.01)	
<b>International application No.</b> PCT/EP00/05152	<b>Applicant's or agent's file reference</b> L/XG34/cm/5
<b>International filing date (day/month/year)</b> 05 June 2000 (05.06.00)	<b>Priority date (day/month/year)</b> 04 June 1999 (04.06.99)
<b>Applicant</b> MARTINEZ-FORCE, Enrique et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

18 December 2000 (18.12.00)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<b>The International Bureau of WIPO</b> 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Juan Cruz Telephone No.: (41-22) 338.83.38
--	---

# PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>L/XG34/cm/5</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/EP 00/05152</b>	International filing date (day/month/year) <b>05/06/2000</b>	(Earliest) Priority Date (day/month/year) <b>04/06/1999</b>
Applicant  <b>CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

### 1. Basis of the report

a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing:

☐ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.

## INTERNATIONAL SEARCH REPORT

International Application No

PC 00/05152

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01H5/10 A23D7/00 A61K7/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01H A61K A23D C11C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

FSTA, BIOSIS, EPO-Internal, COMPENDEX, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 38 31 516 A (HENKEL KGAA) 22 March 1990 (1990-03-22) tables 2,3 ---	1-5, 10-12
A	DATABASE FSTA 'Online! INTERNATIONAL FOOD INFORMATION SERVICE (IFIS), FRANKFURT/MAIN, DE; AN: 1998-09-n0430, 1998 RUÍZ-GUTIÉRREZ, V. ET AL: "Compositions of human VLDL triacylglycerols after ingestion of olive oil and high oleic sunflower oil" XP002148856 & Journal of Nutrition, Vol 128, num. 3, pages 570-576 abstract ----- -/-	1-9

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\* & \* document member of the same patent family

Date of the actual completion of the international search

29 September 2000

Date of mailing of the international search report

13/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fonts Cavestany, A

# INTERNATIONAL SEARCH REPORT

International Application No

PC 00/05152

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>US 5 558 871 A (GRIAT JACQUELINE ET AL)  24 September 1996 (1996-09-24)  claims 4,12</p> <p>-----</p>	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No


PCT/JP 00/05152

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
DE 3831516	A	22-03-1990	NONE		
US 5558871	A	24-09-1996	FR	2714831 A	13-07-1995
			AT	141787 T	15-09-1996
			BR	9500130 A	03-10-1995
			CN	1109741 A	11-10-1995
			DE	69400430 D	02-10-1996
			DE	69400430 T	16-01-1997
			EP	0662318 A	12-07-1995
			ES	2094033 T	01-01-1997
			HU	70586 A, B	30-10-1995
			JP	2648462 B	27-08-1997
			JP	8034708 A	06-02-1996
			PL	306587 A	24-07-1995
			RU	2125982 C	10-02-1999

# PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference L/XG34/ems/5		<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/EP00/05152	International filing date (day/month/year) 05/06/2000	Priority date (day/month/year) 04/06/1999	
International Patent Classification (IPC) or national classification and IPC A01H5/10			
Applicant CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS			
<p>1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.</p> <p>2. This REPORT consists of a total of 4 sheets, including this cover sheet.</p> <p><input type="checkbox"/> This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).</p> <p>These annexes consist of a total of sheets.</p>			
<p>3. This report contains indications relating to the following items:</p> <ul style="list-style-type: none"> <li>I <input checked="" type="checkbox"/> Basis of the report</li> <li>II <input type="checkbox"/> Priority</li> <li>III <input type="checkbox"/> Non-establishment of opinion with regard to novelty, inventive step and industrial applicability</li> <li>IV <input type="checkbox"/> Lack of unity of invention</li> <li>V <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement</li> <li>VI <input type="checkbox"/> Certain documents cited</li> <li>VII <input type="checkbox"/> Certain defects in the international application</li> <li>VIII <input type="checkbox"/> Certain observations on the international application</li> </ul>			
Date of submission of the demand  18/12/2000		Date of completion of this report  05.09.2001	
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465		Authorized officer  Van Woensel, G  Telephone No. +49 89 2399 2089	



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/05152

## I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

**Description, pages:**

1-26 as originally filed

**Claims, No.:**

1-12 as originally filed

**Drawings, sheets:**

1/5-5/5 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/05152

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes:	Claims	1-5
	No:	Claims	6-12
Inventive step (IS)	Yes:	Claims	1-5
	No:	Claims	
Industrial applicability (IA)	Yes:	Claims	1-12
	No:	Claims	

### 2. Citations and explanations **see separate sheet**



**Ad V**

1. The subject-matter of present claim 1 appears to meet the requirements of Article 33 PCT. None of the cited prior art documents discloses an oil and its use comprising a high oleic acid content and a high stearic acid content wherein less than 10%wt of the fatty acid groups in the sn-2 position of the TAG molecules are saturated fatty acid groups.  
Claims 2-5 are dependent on claim 1 and as such also meet the requirements of the PCT with respect to novelty and inventive step.
2. The present application does not meet the requirements of Article 33 PCT, because the subject-matter of claims 6-12 is not new in the sense of Article 33(2) PCT.  
The products of claims 6-12 are defined by one specific oil used. It is not discernable on a sauce for instance, which mixture of oils, fatty acids and other ingredients has been used, because the weight percentages of the respective fatty acids in the end product, will vary from the weight percentage in one of the ingredients.  
Therefore, it is not possible to distinguish the present products from food products or cosmetic products known in the art.
3. Claims 1-12 are considered to be industrially applicable (Article 33(4) PCT).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 December 2000 (14.12.2000)

PCT

(10) International Publication Number  
**WO 00/74471 A1**

- (51) International Patent Classification<sup>7</sup>: A01H 5/10, de la Concepción, E-41907 Valencia de la Concepción (ES).  
A23D 7/00, A61K 7/00
- (21) International Application Number: PCT/EP00/05152
- (22) International Filing Date: 5 June 2000 (05.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/326,501 4 June 1999 (04.06.1999) US  
99204384.4 17 December 1999 (17.12.1999) EP  
60/180,455 4 February 2000 (04.02.2000) US
- (71) Applicant (for all designated States except US): CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS [ES/ES]; Instituto de la Grasa, Avenida Padre García Tejero, 4, E-41012 Sevilla (ES).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MARTINEZ-FORCE, Enrique [ES/ES]; Lamarque de Novoa, 9, 3<sup>a</sup>A, E-41008 Sevilla (ES). MUÑOZ-RUZ, Juan [ES/ES]; Avenida Barcelona, 18, 5<sup>o</sup>-2, E-14010 Córdoba (ES). FERNANDEZ-MARTINEZ, José, María [ES/ES]; Calle Teruel, 10, 3<sup>o</sup>-3, E-14012 Córdoba (ES). GARCES, Rafael [ES/ES]; Calle Dinamarca, 9, Valencia
- (74) Agents: VAN SOMEREN, Petronella, Francisca, Hendrika, Maria et al.; Arnold & Siedsma, Sweelinckplein 1, NL-2517 GK The Hague (NL).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— With international search report.  
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF HIGH OLEIC HIGH STEARIC OILS

(57) Abstract: The invention relates to the use of an oil having an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of said oil, and wherein a maximum of 10 wt% of the fatty acid groups in the sn-2 position of the TAG molecules constituting the oil are saturated fatty acid groups in food and cosmetic products.

WO 00/74471 A1

## USE OF HIGH OLEIC HIGH STEARIC OILS

The present invention relates to the use of an oil having a high oleic and high stearic content in 5 various products.

The uses of oils are determined by their fatty acid composition. The principal component of oils are the triacylglycerol (TAG) molecules, which constitute normally more than 95% of the oil. Three fatty acids are 10 bound to a molecule of glycerol to make the TAG. If these fatty acids are mainly saturated fatty acids ("saturates") the product is called fat and it is solid at room temperature. On the other hand if the fatty acids are mainly unsaturated then it is called oil and it is 15 liquid at room temperature.

The oils obtained from seeds cultivated in temperate climate (sunflower, soybean, rapeseed, etc.) have mainly unsaturated fatty acids, like linoleic and oleic acids, so they are liquid and primarily used for 20 cooking, salad dressing, etc. Fats are obtained from animals (margarine, lard, etc.), some tropical trees (cocoa, palm) or chemically modified (hydrogenation and transesterification) liquid vegetable oils. They have mainly saturated (palmitic or stearic acids) or 25 chemically modified fatty acids (trans fatty acids) all with high melting point.

Table 1 shows as an example the fatty acid composition and other properties of some fats and oils. The fats are needed for most of the food industry to make 30 margarine, shortening, bakery, confectionery, snacks, etc. The food industry uses the fat for these purposes because of their plastic properties (they do not melt, can be spread, or do not stick to the hand) and stability (they have a good resistance to oxidation at room or high 35 temperatures).

Table 1

Oil or fat	Fatty acid composition (%)						Properties	
	Others <sup>1</sup>	Myristic	Palmitic	Stearic	Oleic	Linoleic	Trans	Saturated
5 Lard	3	2	25	12	45	10	1	79
Butter	14	10	26	12	28	3	3	84
Margarine			10	7	46	34	23	*
Palm oil		1	45	5	39	9		18
Olive oil	1		14	3	71	10		2
10 Cocoa butter			26	35	35	3		4
Normal sunflower			7	5	30	57		1
15 High oleic sunflower			5	4	88	2		1

<sup>1</sup> "others" are palmitoleic in the case of lard and olive oil and also fatty acids shorter than 12 carbons in butter

\* depends on the level of hydrogenation

The actual available fats are however not a good option because they have negative nutritional properties. The main problem is that they raise the bad form of serum cholesterol (low density lipoprotein, LDL). This is due to several facts, some related to the origin of the fat and others with the manipulation thereof. Animal fats have most of the saturated fatty acids in the position 2 of the TAG molecule. Most vegetable fats and oils, however, have only minor amounts of saturated fatty acids in this position and are therefore more healthy.

During digestion the TAG molecule is hydrolysed by enzymes called lipases (figure 1). The fatty acids in positions 1 and 3 are liberated as free fatty acids. If these fatty acids are saturated they form insoluble salts with calcium and magnesium, being mostly excreted. But fatty acids in position 2 form with the glycerol a molecule of monoacylglycerol, which has detergent properties and is easily absorbed into the body. The

saturated fatty acids from animal fats are then absorbed, thus raising LDL.

In order to increase the percentage of saturated fatty acids, vegetable oils are hydrogenated and/or transesterified. The hydrogenation process produces trans fatty acids that probably are even worse than saturated fatty acids as illustrated by Willett, W.C. & Ascherio, A. (1994) Trans fatty acids: Are the effects only marginal? American Journal of Public Health 84:722-724. The transesterification process changes randomly the fatty acids within the three positions, converting a healthy vegetable oil with low saturated fatty acid in the 2 position in an oil that has near 30% of saturated fatty acids. So neither of the two chemical modifications leads to a healthy product.

However, not all fats are unhealthy. It has been demonstrated that cocoa butter, which has around 60% of saturated fatty acids, the rest being mainly oleic acid, does not raise serum cholesterol. This is due to two main reasons. One is that only 4% of the saturated fatty acids are in position 2 and the other is that the principal saturated fatty acid is stearic acid. Stearic acid does not have a negative effect on serum cholesterol. Probably the amount of 35% of oleic acid in the cocoa butter also adds to its healthy property.

It is important to note that except in cocoa butter, palmitic acid is the main saturated fatty acid of commodity fats. Palmitic is however not a very healthy fat.

Traditional breeding and mutagenesis has not been the only tool used to form seeds producing oil with different fatty acid profiles. Increases in stearic acid in oil bearing plants have also been addressed by the introduction of transgenes into the germplasm, to alter the fatty acid biosynthesis pathway of the vegetable oil. The fatty acid biosynthesis in vegetable oil, but more particularly sunflower oil, includes the biosynthesis of basically two saturates (palmitate, stearate) and two

unsaturates (oleate and linoleate). In oilseeds, the  
stearoyl-ACP desaturase is the enzyme which introduces  
the first double bond on stearoyl-ACP to form oleoyl-ACP.  
Thus, this is an enzyme that assists in the determination  
5 of the unsaturation in the C18 length fatty acids.

In U.S. Patent No. 5,443,974 the inhibition of  
canola enzyme stearoyl-ACP desaturase was described. The  
stearate levels were increased but the levels of  
palmitate were basically unaffected. Inhibition of the  
10 plant enzyme stearoyl-ACP desaturase in canola was also  
reported by Knutzon et al., Proc. Natl. Acad. Sci. USA  
89:2624-28 (1992). These results showed an increase in  
the level of stearate produced in the canola seed. The  
research also showed that inhibition by antisense in  
15 seeds of canola and soybean, respectively, showed  
increased stearate. When a plasmid containing a gene  
encoding for stearoyl-ACP desaturase was placed in  
canola, this inhibition resulted in an increase in  
stearic acid but unfortunately a reduction in the oleate.  
20 However, in the soybean this inhibition of stearate  
resulted in a less dramatic reduction of the oleate. This  
slower decrease in oleate however may have been a  
function of the small initial levels of oleate in the  
soybean. The fatty acid pathway in most oilseed plants  
25 appears to be resistant to maintaining both oleic and  
stearic at elevated levels.

PCT/US97/01419 describes increased levels of  
both stearic acid and palmitic acid in sunflowers through  
the inhibition of the plant enzyme stearoyl-ACP  
30 desaturase. As indicated above, palmitic oil is not,  
however, viewed as being a very healthy oil.

PCT/US96/09486 discloses that sunflower oil  
levels of both palmitic and oleic acids could be  
increased, the seeds having increased levels of palmitic  
35 acid of 21-23% and of oleic acid of 61%. The sunflower  
oil is liquid at room temperature. But the increased  
palmitic fatty acid level is alleged to allow the oil to  
be used in shortening and in margarine with relatively

low level of hydrogenation, which leads to a relatively low level of trans-fatty acids in the resulting product. However, the commercial value may be questioned because of the high level of palmitic acid.

5           There thus remains a need for a sunflower oil which is both healthy and useful for industrial purposes. Furthermore, it is desirable to have a sunflower oil that has a balance of good saturates and good unsaturates, i.e. that is high in unsaturates but has sufficient  
10 saturates to be used for margarines or hardstock without high levels of hydrogenation, thus leading to no trans-fatty acids in the resulting product. Basically, there remains a need for a sunflower plant that can produce seed containing oil which is high in oleic acid and in  
15 stearic acid with reduced linoleic levels.

          It is therefore the object of the present invention to provide a vegetable oil with high stearic acid (as saturated fatty acid) and high oleic acid (as unsaturated fatty acid) contents that will reduce the  
20 above described problems with fat. In this oil the stearic acid should preferably be in positions 1 and 3 of TAG.

          The present invention is based on the following considerations. The seed fatty acid biosynthesis occurs  
25 inside the plastid (figure 2). A series of cycling reactions catalysed by the enzymatic complex FAS I produces the palmitoyl-ACP that has 16 carbons. A second enzymatic complex called FAS II elongates the palmitoyl-ACP to stearoyl-ACP (18 carbons), that is further modified by  
30 the stearate desaturase to produce oleoyl-ACP. These are the three main fatty acids synthesised by the plastid, being cleaved off the ACP by the action of the enzyme thioesterase and then exported out of the plastid. Later in the cytoplasm, the oleic acid may be desaturated to  
35 linoleic and linolenic acids.

          The TAG (storage oil) is produced in the cytoplasm using the pool of fatty acids in the cytoplasm. This fatty acid pool consists of the fatty acids exported

from the plastid and the linoleic acid made in the cytoplasm by desaturation. Thus, the fatty acid composition of TAG is determined by the fatty acids exported out of the plastid plus the linoleic acid  
5 produced in the cytoplasm.

It was then contemplated that a new plant that is rich in stearic and oleic acids could be selected if a reduced stearate desaturase activity (leading to a decrease in the amount of oleoyl-ACP formed and therefore  
10 in an increase in the stearyl-ACP) was combined with a good thioesterase activity on stearyl-ACP (which leads to the stearic acid being transported out of the plastid into the cytoplasm). This plant will produce an accumulation of stearyl-ACP inside the plastid, and the good  
15 activity of the thioesterase over stearyl-ACP should export it very well out of the plastid, having there a high stearic acid content available for TAG biosynthesis.

Out of the plastid, in the cytoplasm the high oleic character is necessary to keep the linoleic acid  
20 content low. In high oleic lines, the conversion pathway does not work properly, so there is no conversion of oleic acid to linoleic acid.

The present invention is thus based on the finding that by selection of one parent line that has a  
25 high stearic (HS) acid content on the one hand and a second parent line having a high oleic and high thioesterase (HOHT) activity over stearyl-ACP on the other hand, crosses can be made that result in seeds having a combination of the high stearic and high oleic  
30 properties (HSHO). In addition, it was surprisingly found that in said oil a maximum of 10 wt% of the fatty acid groups in the sn-2 position of the TAG molecules are saturated fatty acid groups.

Therefore, the present invention relates to  
35 plant seeds that contain an oil comprising an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of said oil, and wherein a maximum of 10 wt% of the fatty



acid groups in the sn-2 position of the TAG molecules constituting the oil are saturated fatty acid groups. Preferably, the saturated fatty acid groups are stearic acid groups. It is preferred that the oil has in the sn-2 position of the TAG molecules a maximum of 8%, more preferably a maximum of 5 wt% of saturated fatty acid groups, in particular stearic acid groups.

Regarding the other fatty acids, it is preferred that the oleic acid content is from 55 to 75 wt%, the stearic acid content is from 15 to 50 wt%, in particular 20 to 40 wt%, and the linoleic acid content is less than 20 wt%. Preferably the total level of saturated fatty acids is at least 20 wt%.

Selection of the parents can be achieved as follows.

Lines with high stearic acid content are lines having a stearic acid content of more than 12%, preferably more than 20%. One example of such a high stearic (HS) parent line, which was selected after mutagenesis and has a stearic acid content of 26 wt%, is available as "CAS-3" (ATCC deposit no. 75968, deposited on December 14, 1994). Another example is "CAS-4", having a stearic acid content of 16.1 wt% (ATCC deposit no. 75969, deposited on December 14, 1994). By analysing the fatty acid composition of oil derived from the seeds of other candidate lines, the skilled person will be able to select other suitable parent lines.

It was found that some of the usual high oleic varieties could not be used for the purpose of the invention because they were found to have very low thioesterase activity over the stearyl-ACP. To overcome this, by measuring the thioesterase activity, lines with good activity over stearyl-ACP can be selected from the available high oleic lines collections.

In short, one would first analyse the fatty acid composition of the oil of several promising lines. A suitable HOHT parent line would have more than 7-8% stearic acid and either less than 5% linoleic acid or

more than 75% oleic acid. Subsequently, the selected lines must be grown and self pollinated. The total thioesterase activity is measured in seeds 15 days after flowering (15DAF) on both oleoyl-ACP and stearoyl-ACP. In 5 suitable lines, the activity over stearoyl-ACP should be more than 10% of the activity over oleoyl-ACP. The ratio between both activities determines whether a line is suitable as a parent line or not.

In Table 2 the fatty acid composition and 10 thioesterase activity of two high oleic sunflower lines are illustrated.

Table 2

Stearic acid content and thioesterase Vmax over the 15 stearoyl-ACP of 15 days after flowering seeds from two high oleic sunflower lines.

Sunflower line	Stearic acid (%)	Thioesterase activity Vmax
HOHT	17.8	2.03
HOLT	8.0	0.82

20 The HOHT line is a high oleic line with thioesterase over stearoyl-ACP activity (HOHT) of more than twice the thioesterase Vmax over stearoyl-ACP than an usual high oleic line (HOLT). The relative activity of the enzymes 25 over the stearoyl-ACP standardised with respect to the one over oleoyl-ACP is illustrated in Figure 3. This line has as a consequence more stearic acid at 15 days after flowering (Table 2) and also in the oil obtained from the mature seed (Table 3).

30

Table 3

Fatty acid composition (%) of seeds from two high oleic sunflower lines.

35

	Fatty acid composition (%)
--	----------------------------

Sunflower line	palmitic	stearic	oleic	linoleic	araquic	behenic
HOHT	4.3	9.7	78.5	3.9	1.0	2.6
HOLT	3.8	4.9	84.3	4.8	0.5	1.7

5

This HOHT parent line was deposited on September 7, 1999 with the American Type Culture Collection (10801 University Boulevard, Manassas, Va 20110-2209) and was assigned the number PTA-628.

10 Lines of both types (HOHT and HOLT) have been crossed with the high stearic CAS-3 line. In Figures 4 (for HOHT) and 5 (for HOLT), the F2 segregation for both traits (high stearic acid content and high oleic acid content) are shown. The seeds with higher content in  
 15 stearic and oleic acids are within a circle. From the figures it follows that the HOHT line with high thioesterase activity over stearyl-ACP has high oleic high stearic seeds and the line without high thioesterase activity has no seeds of this type. Table 4 shows the  
 20 fatty acid composition of these lines.

Table 4

Fatty acid composition of selected high oleic and stearic  
 25 lines, with high and low thioesterase activity over stearyl-ACP, after crossing with HS line CAS-3

	Fatty acid composition (%)					
Sunflower line	palmitic	stearic	oleic	linoleic	araquic	behenic
30 HOHTxCAS-3	5.2	24.6	59.2	6.8	1.8	2.4
HOLTxCAS-3	4.3	17.4	72.1	4.0	1.3	2.8

The selected F2 lines are selfed for 5 to 6 generations in isolated conditions to avoid  
 35 contamination. The resultant generations are selected,

based on high oleic and stearic acid content.

Thioesterase activity can be analysed to assist in the selection process. Likewise, marker assisted breeding can be employed to track any or all of the three traits to make the selection process quicker. Various markers such as SSR microsatellite, ASO, RFLP and likewise can be employed. The use of markers is not necessary, as standard tests are known for determining oleic, stearic, and thioesterase activity. However, once identified markers make trait tracking easier and earlier in the plant's life.

The true breeding plants produce an oil having a similar fatty acid composition to the F2 seeds selected with a low content of saturated fatty acid in the 2 position of the TAG molecule (Table 5).

Table 5

Fatty acid composition of oil, TAG and sn-positions of true breeding HSHO plants selected. n.d.= not detected.

20

Fatty acid composition (mol%)						
	Palmitic	Stearic	Oleic	Linoleic	Araquic	Behenic
Total oil	5.5	24.9	57.8	8.2	1.7	1.8
TAG	5.6	26.1	57.6	7.4	1.6	1.7
25 sn-2 position	1.7	1.9	87.4	9.0	n.d.	n.d.
sn-1 and 3 position	7.2	33.1	46.8	7.3	2.7	2.9

The invention also relates to plants which form seeds which contain the above described oil of the invention and to the oil per se as well as to products derived from the seeds, such as meal and crushed seeds. The plants, seeds, oil, meal and crushed seeds of the invention are for example sunflower plants, seeds, oil, meal and crushed seeds.

35 The plants and seeds of the invention are obtainable by a method comprising:

a) providing seeds which contain an oil having a stearic acid content of at least 12 wt% based on the total fatty acid content of the oil;

b) providing seeds which contain an oil having an oleic acid content of at least 40 wt% based on the total fatty acid content of the oil, and which have a thioesterase activity over stearyl-ACP that is at least 10% of the thioesterase activity over oleoyl-ACP;

c) crossing plants grown from the seeds provided in step (a) and (b);

d) harvesting the F1 seed progeny.

Preferably, the method further comprises the steps of:

e) planting the F1 progeny seeds to grow plants;

f) self-pollinating the plants thus grown to produce F2 seed;

g) testing the seed for the presence of a stearic acid content in the oil of at least 12 wt% and an oleic acid content of at least 40 wt% and a thioesterase activity over stearyl-ACP that is at least 10% of the thioesterase activity over oleoyl-ACP;

h) planting seeds having the desired levels of stearic acid content, oleic acid content and thioesterase activity to grow plants;

i) self-pollinating the plants thus grown to produce F3 seed; and

j) optionally repeating steps g), h) and i) until the desired levels of stearic acid content, oleic acid content and thioesterase activity are fixed.

Preferably, the stearic acid content is at least 15 wt%, preferably at least 20 wt%.

The present invention also covers the method of obtaining an oil, in particular a sunflower oil, having an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of the oil by extracting oil from the seeds. The method preferably includes an extraction process

which does not involve a substantial modification of the (sunflower) oil.

Additionally, in the process of extraction of the oil from the seeds there is preferably no substantial  
5 chemical or physical modification nor enzymatic rearrangement taking place and preferably no substantial hardening of the oil.

The present invention also includes food products comprising oil obtainable from seeds, in  
10 particular sunflower seeds, having an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of the oil. Food products that are particularly useful for this type of oil include spreads, margarines, shortenings,  
15 sauces, ice-cream, soups, bakery products, confectionery products, and the like. In these food products the level of (sunflower) oil is preferably from 3 to 100 wt% relative to the total oil weight in the product. When used to form a spread according to the present invention  
20 the (sunflower) oil is preferably used as a hardstock at levels of 5 to 20 wt%.

The sunflower seeds of the present invention are also suitable per se for human and animal consumption.

25 The present invention also encompasses cosmetic products comprising an oil, in particular a sunflower oil, the oil having an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of the oil. These  
30 cosmetic products can preferably contain levels of (sunflower) oil from 3 to 100 wt%. Some examples of these cosmetic products would include creams, lotions, lipsticks, soap bars and skin or hair oils.

The present invention also includes a process  
35 for selecting Helianthus annuus plants, capable of producing seeds having the desired oil. The steps of the method are a) selecting a number of Helianthus annuus plants, collecting therefrom the seeds, the oil of which

has a stearic acid content of at least 12 wt% and preferably 18 wt% based on the total fatty acid content; (b) selecting a number of Helianthus annuus plants, collecting therefrom the seeds, which express an oleic acid content of at least 40 wt% based on the oil present in the seed and a thioesterase activity over stearyl-ACP that is at least 10% of the thioesterase activity over oleoyl-ACP; (c) crossing the plants grown from the seeds of (a) and (b); and, harvesting the F1 seed progeny.

Additional steps include the steps of: (d) planting of the seeds or embryo rescue of the embryos of the F1 progeny obtained to form F2 segregating seeds; (e) selecting from the F2 seeds which developed plants, those plants which produce seeds having an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of the oil, optionally selfing the selected plant to form true breeding inbreds.

The present invention also includes the process for producing F1 hybrid seed. The steps of the method are a) planting seed of two inbreds having high oleic acid content of at least 40 wt% and thioesterase activity over stearyl-ACP that is at least 10% of the thioesterase activity over oleoyl-ACP, one of which may be male sterile, b) crossing the two inbreds, and c) harvesting the F1 seed capable of producing F2 seed with an at least 40 wt% oleic acid content and an at least 12 wt% stearic acid content.

The present invention encompasses a vegetable oil with a new and unique fatty acid composition produced in easy to grow crops. The preferred crop is sunflower. This plant was used for making this invention. However, the invention is more broadly applicable and selection of suitable parents to produce the derived vegetable oil could likewise modify other crops. These crops would include at least Brassicas, peanuts, palms and other oil producing plants. When mutation is used for making one or both of the parents, the crop should be susceptible to

mutagenically induced oil changes. Rape seed meets all these requirements as does sunflower, these crops are presently some of the most useful crops for production of this new and unique fatty acid composition in the oil of  
5 their seeds.

In this application reference is made to the following figures:

Figure 1: hydrolysis of triacylglycerols by lipase;

10 Figure 2: plastid showing the fatty acid biosynthesis in oilseeds;

Figure 3: elevated thioesterase activity shown as the relative activity of the thioesterase over stearoyl-ACP and oleoyl-ACP of HOHT and HOLT;

15 Figure 4: the F2 segregation for stearic and oleic acids of the cross between high oleic with high thioesterase activity over stearoyl-ACP line (HOHT) and a high stearic acid line (CAS-3);

20 Figure 5: the F2 segregation for stearic and oleic acids of the cross between high oleic with low thioesterase activity over stearoyl-ACP line (HOLT) and a high stearic acid line (CAS-3).

## 25 DEFINITIONS

"SUNFLOWER" shall mean Helianthus annuus.

"PLANT" shall include the complete plant and all plant and cell parts including pollen, kernel, oil, embryo, stalk, head, roots, cells, meristems, ovule,  
30 anthers, microspores, embryos, DNA, RNA, petals, seeds, and the like and protoplasts, callus or suspensions of any of the above.

"15DAF" shall mean 15 days after flowering.

35 "TOTAL FATTY ACID CONTENT" of the sunflower oil refers to the sum of C16:0, 18:0, 18:1, 18:2, 20:0, 22:0 and the traces of other like fatty acids as determined simultaneously in the oil from the seed.



"HOLT" shall mean having high to medium-high (40%-90%) oleic acid levels in the oil when compared to normal, wildtype sunflower seed (oleic acid levels of 17%-20%) wherein there are "LOW LEVELS OF THIOESTERASE ACTIVITY". A "HOLT LINE" is a line, in particular a sunflower line, having the HOLT trait.

"HOHT" shall mean having high to medium-high (40%-90%) oleic acid levels in the oil when compared to normal, wildtype sunflower seed (oleic acid levels of 17%-20%) wherein there are "HIGH LEVELS OF THIOESTERASE ACTIVITY". A "HOHT LINE" is a line, in particular a sunflower line, that has the HOHT trait.

"HIGH LEVELS OF THIOESTERASE ACTIVITY" shall mean levels (at 15DAF) of thioesterase activity over stearyl-ACP which are at least 10% of the thioesterase activity over oleoyl-ACP. Consequently, "LOW LEVELS OF THIOESTERASE ACTIVITY" shall mean levels which are below the "HIGH LEVELS OF THIOESTERASE ACTIVITY".

"HS" shall mean having stearic acid levels in the oil of at least 12 wt% and preferably at least 15 wt% or more preferably at least 18 wt% or even at least 20 wt% based on the total fatty acid content. "HIGH STEARIC LINE" or "HS LINE" shall mean a line, in particular a sunflower line, having the HS trait.

"HOHS" shall mean having levels of above 40% oleic acid and at least 12 wt% stearic acid in the oil and preferably having levels of at least 15% wt, more preferably at least 18 wt% or even at least 20 wt% stearic acid in the oil. A "HOHS LINE" shall mean a line having the HOHS trait.

#### EXAMPLES

##### INTRODUCTION

##### 35 Preparation of HS parent

In order to obtain the HS parent a method can be used for preparing sunflower seeds having an increased stearic acid and oleic acid content as compared to wild

type seeds. This method includes the step of treating parent seeds with a mutagenic agent during a period of time and in a concentration sufficient to induce one or more mutations in the genetic trait involved in stearic acid or oleic acid biosynthesis. This results in an increased production of stearic acid and/or an increased level of oleic acid. These mutagenic agents include agents such as sodium azide or an alkylating agent, like ethyl methane sulfonate, of course any other mutagenic agent having the same or similar effects may also be used. The treated seeds will contain inheritable genetic changes. These mutated seeds are then germinated and progeny plants are developed therefrom. To increase the traits in the lines the progeny can be crossed or selfed. The progeny seeds are collected and analysed.

Sodium azide and ethyl methane sulfonate were used as mutagenic agents in Example 1. Several sunflower lines with a stearic acid content between 12 and 45% have been obtained. In all these cases the original sunflower parent line for the production of the high stearic acid lines used was RDF-1-532 (Sunflower Collection of Instituto de Agricultura Sostenible, CSIC, Cordoba, Spain) that has from 4 to 7% stearic acid content in the seed oil.

25

#### Selecting the HOHT parent

In principle it is sufficient to screen oleic lines for a HOHT phenotype and use this line for either transformation or for crossing to a high stearic line to develop a HOHS line. A suitable line is at least the HOHT parent line that was deposited on September 7, 1999 with the American Type Culture Collection (10801 University Boulevard, Manassas, Va 20110-2209) and was assigned the number PTA-628.

35

#### Making the HOHS line

Seeds having the HOHT trait or the stearic trait can then be crossed to each other to form the HOHS

line. Optionally there can be additional cycles of germination, culturing, and selfing to fix the homozygosity of the traits in the lines and crossing and collection of seeds.

5

## MATERIALS AND METHODS

### Plants growth conditions

Sunflower (Helianthus annuus L.) seeds from high oleic lines with altered seed fatty acid content was  
10 used to test for the thioesterase activities over stearyl-ACP. Plants were cultivated in growth chambers at 25/15°C (day/night) temperature, 16 hours photoperiod and photon flux density of 300 micromol m<sup>-2</sup>s<sup>-1</sup>. Seeds for analysis were harvested at 15 days after flowering and  
15 kept at -20°C.

### Radioactive reagents and preparation of acyl-ACPs

1-<sup>14</sup>C-Oleic with specific radioactivity of 2.1 GBq/mmol and [9,10(n)-<sup>3</sup>H] stearic acid with specific  
20 radioactivity of 1.9 GBq/mmol were obtained from American Radiolabeled Chemicals Inc. (St.Louis, Mo., USA). To prepare the fatty acid sodium salt, an appropriate volume of fatty acid solution was transferred to a glass tube, the solvent was removed under a stream of nitrogen, and  
25 the residue was dissolved in 10% Triton X-100, 0.6 mM NaOH. This solution was heated at 55°C for 1 hour to ensure homogeneity.

Acyl-ACPs were prepared using a modification of the enzymatic synthesis procedure of Rock C.O. et al.  
30 (1981) Methods Enzymology 72:397-403. Assays contained 0.1 M Tris-HCl (pH 8.0), 0.4 M LiCl, 5 mM ATP, 10 mM MgCl<sub>2</sub>, 2 mM DTT, 130 microM fatty acid sodium salt, 0.27 mM ACP-SH and 1.8 mU of acyl-ACP synthetase (the last two components were purchased from Sigma-Aldrich Quimica S.A.  
35 Madrid, Spain) in a final volume of 110 microliter. Reactions were incubated at 37°C for 3 hours. After this time the pH was acidified to 6.0 by adding 1 microliter of 3.6 M HCl and the mixture was cleaned of free fatty

acids using a modification of the method described by Mancha M. et al. ((1975) Anal. Biochem. 68:600-608), which method consists of adding an equal volume of isopropanol and washing three times with hexane saturated 5 in water/isopropanol (1:1; v/v).

Preparation of crude extracts for enzyme assays and protein determination

Frozen seeds were peeled and ground in extract 10 buffer containing 20 mM Tris-HCl (pH 8.5), 2 mM DTT and 5% (v/v) glycerol (Dörmann P. et al. (1994) Biochim. Biophys. Acta 1212:134-136) at 1 g of tissues/10 ml of buffer. Protein concentrations were measured using a Protein Assay Kit (Bio-Rad) according to the 15 manufacturer's recommendations, with BSA as standard.

Enzyme assays

Acyl-ACP thioesterase activity was assayed in a final volume of 170 microliter using 130 microliter of 20 crude extract. Control assays had crude extract omitted. Reactions mixtures contained 20 mM Tris-HCl (pH 8.5), 5% glycerol and 2 mM dithiothreitol (DTT) and different concentrations of substrates (stearoyl-ACP and oleoyl-ACP). Incubations were carried out for 20 min at 25°C. 25 Reactions were stopped by the addition of 170 microliter of 1 M acetic acid in isopropanol containing 1 mM of oleic acid. Mixtures were then washed three times with hexane saturated in water/isopropanol (1:1, v/v).

Acyl-ACP thioesterase activity was determined 30 by counting the radioactivity of the aqueous phase, which contained the non-hydrolysed substrates. Then, 3 ml of solvent scintillant (purchased from National Diagnostics, Hesse, England) was added and the radioactivity was measured using a scintillation counter (Rackbeta II; LKB, 35 Sweden). Data from acyl-ACP thioesterase assays were fitted to the Michaelis-Menten equation by non-linear least-squares regression analysis using Microcal Origin 4. 1, and correlated to  $P < 0.05$ , as determined by paired

Student's test. Vmax and Km were derived from these curves.

#### EXAMPLE 1

##### 5 Preparation of a HS line

###### 1. Mutation with EMS

Seeds were mutagenised with a solution of 70 mM of ethyl methane sulfonate (EMS) in water. The treatment was performed at room temperature during 2 hours while  
10 shaking (60 rpm). After mutagenesis the EMS solution was discarded and seeds were washed during 16 hours under tap water.

Treated seeds were germinated in the field and plants were self-pollinated. The seeds collected from  
15 these plants were used to select new sunflower lines with modifications in the fatty acid composition. By using the method of Garcés, R. and Mancha, M. ((1993) Anal. Biochem. 211, 139-143) the seed fatty acid composition was determined by gas liquid chromatography, after  
20 converting the fatty acids into their corresponding methyl esters.

A first plant with 9 to 17% stearic acid content in the oil was selected. The progeny was cultivated for five generations wherein the stearic acid  
25 content increased and the new genetic trait became stably fixed in the genetic material of the seed. This line is called CAS-3. The minimum and the maximum stearic acid content of the line were 19 and 35% respectively. The stearic acid content of oil extracted from seeds from  
30 this cell line may thus lie between 19 and 35%.

###### 2. Mutation with sodium azide

Sunflower seeds were mutagenised with sodium azide, at a concentration of 2 mM in water. The treatment  
35 was performed at room temperature during two hours while shaking (60 rpm). Then the mutagenesis solution was discarded and seeds were washed during 16 hours with tap water.

Seeds were planted in the field and plants were self-pollinated. Seeds from these plants were collected, and the fatty acid composition was determined by gas liquid chromatography, after converting the fatty acids into their corresponding methyl esters using the method described in Example 1.

Seeds from a plant having around 10% stearic acid in the oil were selected and cultivated for five generations. During this procedure the stearic acid content was increased and the new genetic trait fixed. This line is called CAS-4. A selected sample of this line was analysed resulting in a stearic acid content of 16.1%. The minimum and the maximum values were 12 and 19%, respectively.

15

Table 6

Percentage fatty acids				
Line	Palmitic	Stearic	Oleic	Linoleic
20 CAS-3	5.1	26.0	13.8	55.1
CAS-4	5.5	16.1	24.3	54.1

CAS-3 and CAS-4 are on deposit with the American Type Culture Collection, having ATCC numbers 75968 and 75969, respectively.

**EXAMPLE 2**Production of a HSHO line30 1. General

Sunflower plants were grown from the sunflower seeds of the HOHT line, seeds of which are on deposited at ATCC (PTA-628). Sunflower plants were also grown from the sunflower seeds of CAS-3. The lines were crossed. The plants were assisted by artificial pollination in order to ensure adequate seed production occurred. The F1 seed was produced on the HOHT line, or vice versa, and harvested. The F2 seeds with more than 20% stearate and

more than 40% oleate were selected. Although this produces the oil of the present invention the level of production is limited.

Therefore fixed inbred lines evidencing seeds  
5 with these oil profiles are desirable. These homozygous fixed inbred HSHO lines can then be crossed to form hybrid seed, which will produce F2 seed evidencing the desired oil traits of the present invention.

Toward this end the F1 seeds were planted and  
10 produced plants were selfed in isolated conditions and F2 seed was produced. The F2 seed was tested for the three traits, high stearic, high oleic and high levels of thioesterase activity. The remaining portion of the seeds evidencing these traits was employed to grow plants to  
15 form F3 seed. The selfing and screening and selection process is repeated to develop the fixed homozygous HSHO line, having the following fatty acid profile, C:16 5.4, C:18.0 24.8, C:18.1 58.5, C:18.2 7.2. Once the trait is fixed similar HSHO lines can cross to form hybrid seed  
20 having both traits.

According to the invention sunflower plants and seeds from which said oil can be extracted have been obtained by means of a biotechnological process. This high stearic acid content is an inheritable trait and is  
25 fairly independent from the growing conditions.

## 2. First cross

A sunflower plant was grown from a sunflower seed of an HOHT line having a stearic acid content of  
30 10.7 wt% and an oleic acid content of 74.6 wt%. A sunflower plant was also grown from a CAS-3 sunflower seed. The plants were crossed. The plants were assisted by artificially pollination in order to ensure adequate seed production occurred. The F1 seed was produced on the  
35 HOHT line, or vice versa, and harvested.

A F1 seed having a stearic acid content of 9.8 wt% and an oleic acid content of 80.7 wt%, was selected. This F1 seed was planted and produced a plant which was

selfed in isolated conditions and F2 seeds were produced. These F2 seeds were tested for oleic and stearic acid contents. A seed containing 23.6 wt% of stearic acid and 65.5 wt% of oleic acid was selected.

5           This F2 seed was planted and produced a plant which was selfed in isolated conditions and at 15DAF several seeds were collected and analysed for stearyl-ACP thioesterase activity. Plants with seeds rendering more than 10% stearyl-ACP thioesterase referred to the  
10   oleoyl-ACP thioesterase activity of the same plant were selected.

          Mature seeds from the plants selected in the previous step and having stearic acid content higher than 20 wt% and oleic acid content higher than 40 wt% were  
15   submitted to the selfing, screening and selection process repeatedly to develop the fixed homozygous high stearic high oleic line having the following fatty acid profile in the oil:

          palmitic 7.8 wt%;  
20       stearic 24 wt%;  
          oleic 57.7 wt%;  
          linoleic 5.9 wt%;  
          araquic 1.9 wt%;  
          behenic 2.7 wt%.

25   Once the trait is fixed, similar high stearic high oleic lines can cross to form hybrid seed having the above selected traits.

          An analysis of the sn-2 position and sn-1,3 positions of the TAG molecules of this oil indicates the  
30   following distribution of fatty acids (in wt%):  
      sn-2:

          palmitic 3.3%;  
          stearic 3.4%;  
          oleic 88.8%;  
35       linoleic 4.5%;  
          araquic 0%;  
          behenic 0%

      sn-1,3:



palmitic 9%;  
stearic 29.9%;  
oleic 51.1%;  
linoleic 4.7%;  
5 araquic 2.3%;  
behenic 3%

Thus, the total amount of saturated fatty acid groups in the sn-2 position of the TAG molecules of this oil is 6.7 wt%.

10

### 3. Second cross

A sunflower plant was grown from a sunflower seed of an HOHT line having a stearic acid content of 8.4 wt% and an oleic acid content of 78.5 wt%. A sunflower  
15 plant was also grown from a CAS-3 sunflower seed. The plants were crossed. The plants were assisted by artificially pollination in order to ensure adequate seed production occurred. The F1 seed was produced on the HOHT line, or vice versa, and harvested. A F1 seed having a  
20 stearic acid content of 7.1 wt% and an oleic acid content of 84.6 wt%, was selected. This F1 seed was planted and produced a plant which was selfed in isolated conditions and F2 seeds were produced. These F2 seeds were tested for oleic and stearic acid contents. A seed containing  
25 22.8 wt% of stearic acid and 64.8 wt% of oleic acid was selected.

This F2 seed was planted and produced a plant which was selfed in isolated conditions and at 15 DAF several seeds were collected and analysed for stearoyl-  
30 ACP thioesterase activity. Plants with seeds rendering more than 10% stearoyl-ACP thioesterase referred to the oleoyl-ACP thioesterase activity of the same plant were selected. Mature seeds from the plants selected in the previous step and having stearic acid content higher than  
35 20 wt% and oleic acid content higher than 40 wt% were submitted to the selfing, screening and selection process repeatedly to develop the fixed homozygous high stearic

high oleic line having the following fatty acid profile in the oil:

- 5 palmitic 5.8 wt%;
- stearic 24.7 wt%;
- oleic 57.6 wt%;
- linoleic 8.2 wt%;
- araquic 1.8 wt%;
- behenic 1.9 wt%.

Once the trait is fixed, similar high stearic high oleic lines can cross to form hybrid seed having the above selected traits.

An analysis of the sn-2 position and sn-1,3 positions of the TAG molecules of this oil indicates the following distribution of fatty acids (in wt%):

15 sn-2:

- palmitic 1.7%;
- stearic 1.9%;
- oleic 87.5%;
- linoleic 8.9%;
- 20 araquic 0%;
- behenic 0%

sn-1,3:

- palmitic 7.2%;
- stearic 33.2%;
- 25 oleic 46.9%;
- linoleic 7.3%;
- araquic 2.6%;
- behenic 2.8%.

Thus, the total amount of saturated fatty acid groups in the sn-2 position of the TAG molecules of this oil is 3.6 wt%.

#### 4. Third cross

A sunflower plant was grown from a sunflower seed of an HOHT line having a stearic acid content of 9.9 wt% and an oleic acid content of 81.2 wt%. A sunflower plant was also grown from a CAS-3 sunflower seed. The plants were crossed. The plants were assisted by

artificially pollination in order to ensure adequate seed production occurred. The F1 seed was produced on the HOHT line, or vice versa, and harvested.

A F1 seed having a stearic acid content of 8.9 wt% and an oleic acid content of 82.3 wt%, was selected. This F1 seed was planted and produced a plant which was selfed in isolated conditions and F2 seeds were produced. These F2 seeds were tested for oleic and stearic acid contents. A seed containing 23.9 wt% of stearic acid and 64.0 wt% of oleic acid was selected.

This F2 seed was planted and produced a plant which was selfed in isolated conditions and at 15 DAF several seeds were collected and analysed for stearyl-ACP thioesterase activity. Plants with seeds rendering more than 10% stearyl-ACP thioesterase referred to the oleoyl-ACP thioesterase activity of the same plant were selected. Mature seeds from the plants selected in the previous step and having stearic acid content higher than 20 wt% and oleic acid content higher than 40 wt% were submitted to the selfing, screening and selection process repeatedly to develop the fixed homozygous high stearic high oleic line having the following fatty acid profile in the oil:

palmitic 5.4 wt%;  
stearic 24,2 wt%;  
oleic 62.1 wt%;  
linoleic 4.7 wt%;  
araquic 1.6 wt%;  
behenic 2.0 wt%.

Once the trait is fixed, similar high stearic high oleic lines can cross to form hybrid seed having the above selected traits.

An analysis of the sn-2 position and sn-1,3 positions of the TAG molecules of this oil indicates the following distribution of fatty acids (in wt%):

sn-2:

palmitic 1.8%;  
stearic 3.3%;

oleic 89.6%;  
linoleic 5.3%;  
araquic 0%;  
behenic 0%

5 sn-1,3:

palmitic 9.5%;  
stearic 33.5%;  
oleic 48.2%;  
linoleic 4.3%;

10 araquic 2.2%;  
behenic 2.3%

Thus, the total amount of saturated fatty acid groups in the sn-2 position of the TAG molecules of this oil is 5.1 wt%.

## CLAIMS

1. Use of an oil having an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of said oil, and wherein a maximum of 10 wt% of the fatty acid groups in the sn-2 position of the TAG molecules constituting the oil are saturated fatty acid groups in a food product or cosmetic product.
2. Use as claimed in claim 1, wherein the oil is a sunflower oil.
3. Use as claimed in claim 1 or 2, wherein the food product is selected from the group of spreads, sauces, ice-cream, soups, bakery products and confectionery products.
4. Use as claimed in claims 2 and 3, wherein the food product is a spread in which the sunflower oil is used as a hardstock at a level of 5 to 20 wt%.
5. Use as claimed in claim 1 or 2, wherein the cosmetic product is selected from the group of creams, lotions, lipsticks, soap bars and skin or hair oils.
6. Food product comprising an oil having an oleic acid content of more than 40 wt% and a stearic acid content of more than 12 wt% based on the total fatty acid content of said oil, and wherein a maximum of 10 wt% of the fatty acid groups in the sn-2 position of the TAG molecules constituting the oil are saturated fatty acid groups.
7. Food product as claimed in claim 6, wherein said oil is a sunflower oil.
8. Food product as claimed in claim 6 or 7 selected from the group of spreads, sauces, ice-cream, soups, bakery products and confectionery products.
9. Food product according to claims 6-8, being a spread in which the oil is a sunflower oil and is used as a hardstock at a level of 5 to 20 wt%.
10. Cosmetic product comprising an oil having an oleic acid content of more than 40 wt% and a stearic

acid content of more than 12 wt% based on the total fatty acid content of said oil, and wherein a maximum of 10 wt% of the fatty acid groups in the sn-2 position of the TAG molecules constituting the oil are saturated fatty acid groups.

11. Cosmetic product as claimed in claim 10, wherein the oil is a sunflower oil.

12. Cosmetic product as claimed in claim 10 or 11 selected from the group of creams, lotions, lipsticks, soap bars and skin or hair oils.




Figure 1




Figure 2






Figure 3




Figure4




Figure 5

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/05152

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01H5/10 A23D7/00 A61K7/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01H A61K A23D C11C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

FSTA, BIOSIS, EPO-Internal, COMPENDEX, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 38 31 516 A (HENKEL KGAA) 22 March 1990 (1990-03-22) tables 2,3	1-5, 10-12
A	--- DATABASE FSTA 'Online! INTERNATIONAL FOOD INFORMATION SERVICE (IFIS), FRANKFURT/MAIN, DE; AN: 1998-09-n0430, 1998 RUÍZ-GUTIÉRREZ, V. ET AL: "Compositions of human VLDL triacylglycerols after ingestion of olive oil and high oleic sunflower oil" XP002148856 & Journal of Nutrition, Vol 128, num. 3, pages 570-576 abstract --- -/--	1-9



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

29 September 2000

Date of mailing of the international search report

13/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fonts Cavestany, A

# INTERNATIONAL SEARCH REPORT

Intern. Application No.

PCT/EP 00/05152

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>US 5 558 871 A (GRIAT JACQUELINE ET AL)  24 September 1996 (1996-09-24)  claims 4,12</p> <p>-----</p>	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/EP 00/05152

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3831516 A	22-03-1990	NONE	
US 5558871 A	24-09-1996	FR 2714831 A	13-07-1995
		AT 141787 T	15-09-1996
		BR 9500130 A	03-10-1995
		CN 1109741 A	11-10-1995
		DE 69400430 D	02-10-1996
		DE 69400430 T	16-01-1997
		EP 0662318 A	12-07-1995
		ES 2094033 T	01-01-1997
		HU 70586 A,B	30-10-1995
		JP 2648462 B	27-08-1997
		JP 8034708 A	06-02-1996
		PL 306587 A	24-07-1995
		RU 2125982 C	10-02-1999